

Monte Carlo Simulations of Medium-Scale CMB Anisotropy

A. Kogut^{1,2} and G. Hinshaw¹

Submitted to *The Astrophysical Journal Letters*
January 26, 1996

ABSTRACT

Recent detections of cosmic microwave background (CMB) anisotropy at half-degree angular scales show considerable scatter in the reported amplitude even at similar angular resolution. We use Monte Carlo techniques to simulate the current set of medium-scale CMB observations, including all relevant aspects of sky coverage and measurement technique. The scatter in the reported amplitudes is well within the range expected for the standard cold dark matter (CDM) cosmological model, and results primarily from the restricted sky coverage of each experiment. Within the context of standard CDM current observations of CMB anisotropy support the detection of a “Doppler peak” in the CMB power spectrum consistent with baryon density $0.01 < \Omega_b < 0.13$ (95% confidence) for Hubble constant $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The uncertainties are approximately evenly divided between instrument noise and cosmic variance arising from the limited sky coverage.

Subject headings: cosmic microwave background – nucleosynthesis

¹ Hughes STX Corporation, Laboratory for Astronomy and Solar Physics, Code 685, NASA/GSFC, Greenbelt MD 20771.

² E-mail: kogut@stars.gsfc.nasa.gov.

1 Introduction

The anisotropy of the cosmic microwave background probes the distribution of matter and energy prior to the epoch of structure formation. On large angular scales ($\theta \gtrsim 2^\circ$), CMB anisotropies reflect perturbations larger than the particle horizon and thus probe the primordial density distribution. On angular scales smaller than 2° , causal processes become important and modify the primordial distribution in model-specific ways. Observations of the CMB at angular scales near 0.5° sample these causal processes and offer a powerful test of competing models of structure formation.

The *Cosmic Background Explorer* provides a clean detection of CMB anisotropy on large angular scales, measuring the amplitude of the fluctuations to a significance of 14 standard deviations (Bennett et al. 1996). On smaller angular scales, the situation is less clear. A number of groups have reported detections of CMB anisotropy on angular scales 0.5° to 1.5° (Figure 1). A large scatter is evident, even for experiments at similar angular scales. Does this scatter reflect cosmology (e.g. non-Gaussian features in the distribution of CMB anisotropy), instrumentation, or simply sample variance from the small sky coverage (typically less than 0.1% of the celestial sphere) of an individual measurement?

Scott, Silk, & White (1995) approach this problem analytically, fitting a phenomenological approximation of the CMB power spectrum to recent detections to obtain best-fit shape parameters. They conclude that the data do not show excessive scatter and obtain best-fit parameters consistent with the existence of a “Doppler peak” in the power spectrum at angular scales near 0.5° . In this *Letter* we use Monte Carlo techniques to explore further the issue of scatter between different experiments and the degree to which instrument noise, calibration error, and sky coverage may be expected to affect the results. For specificity, we work in the context of the standard cold dark matter cosmology (density $\Omega_0 = 1$, cosmological constant $\Lambda = 0$, and Hubble constant $H_0 = h \text{ km s}^{-1} \text{ Mpc}^{-1}$ with $h = 0.5$), although our results may be extended to other choices of model parameters.

2 Simulations

We represent the CMB temperature field at angular position (θ, ϕ) as a sum over spherical harmonic amplitudes, $T_{CMB} = \sum_{\ell m} a_{\ell m} Y_{\ell m}(\theta, \phi)$. The amplitudes $a_{\ell m}$ are specified by the cosmological model: for a scale-free power spectrum $P(k) \propto k^n$ they are random Gaussian variables with zero mean and variance

$$\langle a_{\ell m}^2 \rangle = C_\ell = \frac{4\pi}{5} Q^2 \frac{\Gamma[l + (n-1)/2] \Gamma[(9-n)/2]}{\Gamma[l + (5-n)/2] \Gamma[(3+n)/2]}, \quad (1)$$

where Q is the ensemble-averaged quadrupole amplitude (Bond & Efstathiou 1987). The matter content of the universe will modify the coefficients C_ℓ to increase power

at $\ell \sim 200 - 1000$ and produce the so-called ‘‘Doppler peaks’’ in the power spectrum. In this *Letter* we use CDM power spectra (Stompor 1994) normalized to the 4-year *COBE* value $Q = 18 \mu\text{K}$ (Bennett et al. 1996), evaluated over the range $2 \leq \ell \leq 500$. An experiment with Gaussian beam dispersion σ_b will observe a smoothed temperature distribution $T_{\text{obs}}(\theta, \phi) = \sum_{\ell m} a_{\ell m} Y_{\ell m}(\theta, \phi) \exp(-\frac{1}{2}\ell(\ell+1)\sigma_b^2)$. To reduce instrumental artifacts, experimenters rapidly sweep the antenna beam between two or more locations on the sky, and combine these to form temperature differences ΔT whose distribution can then be compared to theory.

A single realization of the CMB sky is specified by the set of amplitudes $a_{\ell m}$. We generate 2000 CMB realizations and use each realization to simulate the current set of medium-scale CMB experiments. With a fixed set of amplitudes $a_{\ell m}$ we evaluate T_{obs} along the scan patterns and with the beam dispersions of the following experiments: ACME/SP94 (Gundersen et al. 1995), ARGO (De Bernardis et al. 1994), IAB (Piccirillo and Calisse 1993), MAX4 γ UMi (Devlin et al. 1994), MAX4 σ Her and ι Dra (Clapp et al. 1994), MSAM 2-beam and 3-beam (Cheng et al. 1994), Python II (Ruhl et al. 1995) and Saskatoon ’94 (Netterfield et al. 1995). For each experiment, we combine the smoothed temperatures T_{obs} to form the appropriate set of noiseless temperature differences ΔT (single difference, double difference, and so forth). We then add random uncorrelated noise to each set of simulated ΔT and multiply the noisy result by a factor $(1 + \delta G)$ where δG is a Gaussian of mean zero and width determined by the calibration uncertainty of each experiment (typically, $|\delta G| \sim 0.1$). We remove the mean value from each noisy simulated experiment and further remove a linear drift for those groups performing such subtraction. To speed processing, we neglect possible foreground emission and correlations from atmospheric or detector noise; in cases where the experimenters observe at multiple frequencies, we use a single frequency with noise appropriate to the final CMB weighted mean used by each group.

From each sky simulation we thus generate a set of simulated CMB experiments, measured at the locations and in a similar manner as the actual experiments. Cosmic variance, instrument noise, and calibration uncertainties are automatically accounted for. We may then compare the scatter of the simulated experiments in each realization to the scatter in the actual data.

We quantify the amplitude of the CMB anisotropy for each simulated experiment using the quadrupole normalization Q_{flat} , the equivalent power for a flat power spectrum (Eq. 1 with $n = 1$) fitted to the simulated temperature differences. For each simulated experiment, we derive Q_{flat} by maximizing the likelihood

$$\mathcal{L}(Q_{\text{flat}}^{ij}) \propto \frac{\exp(-\frac{1}{2} \sum_{\alpha\beta} \Delta T_{\alpha}^{ij} (\mathbf{M}^{-1})_{\alpha\beta}^i \Delta T_{\beta}^{ij})}{\sqrt{\det(\mathbf{M}^i)}},$$

where $\mathbf{M}(Q_{\text{flat}})^i$ is the flat-spectrum covariance matrix between pixels α and β computed from the simulations, the index i tracks the different experiments for each

realization, and j tracks the 2000 CMB realizations. Note that the subtraction of the mean (and sometimes linear drift) from the noisy ΔT correlates the pixels; we account for this by simulating noisy flat-spectrum CMB realizations, computing the mean covariance matrix at different values of Q_{flat} , and inverting the (noisy) matrix $\mathbf{M}$ separately for each Q_{flat} value.

3 Results

For a fixed CDM power spectrum, we generate a set of simulated values Q_{flat}^{ij} for the i th experiment in the j th realization. We then define a χ^2 statistic to quantify the scatter in each realization,

$$\chi_j^2 = \sum_i \left(\frac{(Q_{\text{flat}}^{ij} - \langle Q_{\text{flat}}^i \rangle)}{\delta Q^i} \right)^2,$$

where δQ^i is the standard deviation of each simulated Q_{flat}^i evaluated across all 2000 realizations. The set of 2000 simulated χ^2 values can then be compared to the χ^2 of the actual data calculated in a similar fashion (for a recent tabulation see Scott, Silk, & White 1995).

The actual CMB experiments have $\chi^2 = 10.1$ for 10 degrees of freedom when compared to a CDM model with baryon density fixed at $\Omega_b = 0.05$. 42% of the $\Omega_b = 0.05$ simulations had larger χ^2 , placing the current set of CMB experiments near the median of the simulated distribution. There is no evidence for anomalous scatter in recent detections of CMB anisotropy at half-degree angular scales. We repeat the analysis for models with $0 \leq \Omega_b \leq 0.2$. A flat “no-baryon” universe ($\Omega_0 = 1$, $\Omega_b = 0$, $h = 0.5$) has $\chi^2 = 43.0$ and is rejected at more than 5 standard deviations. CMB anisotropy data clearly support the detection of additional power at half-degree angular scales compared to the *COBE* normalization at large angular scales. In the context of CDM with standard recombination, we minimize χ^2 to obtain a 95% confidence interval for the baryon density $0.01 < \Omega_b < 0.13$. Since we have fixed all other cosmological parameters (curvature, cosmological constant, Hubble constant, recombination), this range for Ω_b can not be regarded as definitive; nevertheless, it is reassuring to find the values derived from CMB anisotropy to be in agreement with standard Big Bang nucleosynthesis (Walker et al. 1991).

To what extent do instrument noise, calibration uncertainty, and limited sky coverage contribute to the uncertainty in recent medium-scale CMB measurements? We repeated the simulations with the same (noiseless) $a_{\ell m}$ coefficients and varied the instrument noise and calibration errors. For noise levels and sky coverage typical of recent medium-scale experiments, the combined uncertainties in Q_{flat} are in the range 6–12 μK , split almost evenly between cosmic variance and instrument noise; calibration uncertainties are generally a minor contribution. Cosmic variance scales as the inverse square root of the sky coverage (Scott, Srednicki, & White 1994;

Hinshaw, Bennett, & Kogut 1994) and places a lower limit to the accuracy achieved at any noise level. Determination of the CMB anisotropy amplitude to 10% accuracy at 0.5° angular scales, let alone multi-parameter fits to the power spectrum, will require observations with significantly greater sky coverage than is currently achieved. Assuming noise levels less than $30 \mu\text{K}$ per field of view, a 10% determination of Q_{flat} requires observing a $30^\circ \times 30^\circ$ patch on the sky.

This work was supported in part by NASA RTOP 399-20-61-01.

References

- Bennett, C.L., et al. 1996, ApJ Letters, submitted
- Bond, J.R., & Efstathiou, G. 1987, MNRAS, 226, 655
- Cheng, E.S., et al. 1994, ApJ, 422, L37
- Clapp, A.C., et al. 1994, ApJ, 433, L57
- De Bernardis, P., et al. 1994, ApJ, 422, L33
- Devlin, M.J., et al. 1994, ApJ Letters, 430, L1
- Gundersen, J.O., et al. 1995, ApJ, 443, L57
- Hinshaw, G., Bennett, C.L., & Kogut, A. 1994, ApJ, 441, L1
- Netterfield, C.B., Jarosik, N., Page, L., Wilkinson, D., & Wollack, E. 1995, ApJ, 445, L69
- Piccirillo, L., & Calisse P., 1993, APJ, 411, 529
- Ruhl, J.E., Dragovan, M., Platt, S.R., Kovac, J., & Novak, G. 1995, ApJ, 453, L1
- Scott, D., Srednicki, M., & White, M. 1994, ApJ, 421, L5
- , Silk, J., & White, M. 1995, Science, 268, 829
- Stompor, R. 1994, A&A, 287, 693
- Walker, T.P., Steigman, G., Schramm, D.N., Olive, K.A., & Kang, H.-S. 1991, ApJ, 376, 51




Figure 1: Recent detections of CMB anisotropy, fitted to the amplitude Q_{flat} of a scale-invariant power spectrum with power equal to the observed power. The grey band indicates the 4-year *COBE* normalization. The solid lines show the predicted power spectrum for standard CDM (upper curve) and $\Omega_b = 0$ (lower line).